

Twenty Years of Debate with Stephen

L. Susskind

Department of Physics
Stanford University
Stanford, CA 94305-4060

Abstract

This is my contribution to Stephen Hawking's 60th birthday party. Happy Birthday Stephen!

1 Crisis and Paradigm Shift

Stephen, as we all know, is by far the most stubborn and infuriating person in the universe. My own scientific relation with him I think can be called adversarial. We have disagreed profoundly about deep issues concerning black holes, information and all that kind of thing. At times he has caused me to pull my hair out in frustration – and you can plainly see the result. I can assure you that when we began to argue more than two decades ago, I had a full head of hair.

I can also say that of all the physicists I have known he has had the strongest influence on me and on my thinking. Just about everything I have thought about since 1980 has in one way or another been a response to his profoundly insightful question about the fate of information that falls into a black hole [1]. While I firmly believe his answer was wrong, the question and his insistence on a convincing answer has forced us to rethink the foundations of physics. The result is a wholly new paradigm that is now taking shape. I am deeply honored to be here to celebrate Stephen's monumental contributions and especially his magnificent stubbornness.

The new paradigm whose broad outlines are already clear involves four closely related ideas which I will call Horizon Complementarity (also known as Black Hole Complementarity) [2, 3], the Holographic Principle [4, 5], the Ultraviolet/Infrared connection [6] and the counting of black hole microstates [7, 8, 9, 10, 11]. Each has had strong support from the mathematics of string theory. We have also learned that certain lessons derived from quantum field theory in a fixed background can lead to totally wrong conclusions. For example quantum field theory gives rise to an ultraviolet divergent result for the entropy in the vicinity of a horizon [12, 13] and a volume's worth of degrees of freedom in a region of space. Another misleading result of quantum field theory is that increasingly large energy and momentum scales are equivalent to progressively smaller distance scales. Finally quantum field theory in a black hole background inevitably leads to the loss of quantum coherence from the vantage point of a distant observer [1].

2 Stephen's Argument for Coherence Loss

Let's begin with the issue of coherence loss. Begin by drawing the Penrose diagram for black hole formation. Let us think of it as a background on which we can study conventional quantum field theory. Now let's add some additional particles which for simplicity are

assumed massless. Incoming particles enter the geometry on past light-like infinity, $\mathcal{I}^-$. The initial Hilbert space of such particles is labeled $\mathcal{H}_-$. The particles interact through Feynman diagrams and some go out to $\mathcal{I}^+$ where they are seen by a distant observer. But some particles end up at the space-like singularity. A conventional local description would have the final states live in a Hilbert space which is a product $\mathcal{H}_+ \otimes \mathcal{H}_{sing}$.

Applying ordinary rules to this system we expect the final state to be given in terms of the initial state by an S matrix,

$$|f\rangle = S|i\rangle \tag{2.1}$$

but to an observer outside the black hole the final state consists only of the particles on $\mathcal{I}^+$. It is therefore described by a density matrix

$$\rho = Tr_{sing}|f\rangle\langle f|. \tag{2.2}$$

Since black holes eventually evaporate we are left with a mixed state and a loss of quantum coherence. I have simplified some of the important parts of Stephen's argument but in the original paper he makes a very convincing case that quantum field theory in a black hole environment leads to a loss of coherence after the evaporation has occurred. The problem with this is that loss of quantum coherence violates the fundamental tenets of quantum mechanics. That of course was Stephen's point.

Historically, for a long time high energy physicists, who had other fish to fry, ignored the problem. By and large, with one or two exceptions [14], the relativity community accepted the loss of coherence without question. However, 't Hooft and I were both deeply disturbed by Stephen's conclusion. In my case I felt that loss of coherence, if it were to occur, could not be quarantined or isolated to phenomena which involved massive black holes. It would infect the rest of physics and cause dramatic disasters in ordinary situations [15]. 't Hooft also expressed similar concerns [16].

There was of course an alternative that I have not mentioned. We might suppose that instead of falling through the horizon, particles (and falling observers) encounter an obstruction, a "brick wall" instead of the horizon. Again the trouble is that an observer, now the freely falling one, would experience a violation of a sacred law of nature, this time the equivalence principle. The horizon according to general relativity is an almost flat region which should behave like empty space, not like a brick wall. It seems that there is no way out; at least one or another observer must witness violations of the usual laws of nature.

At this point one could ask why not just assume neither observer sees a violation of sacred principles. Just assume that the infalling information is “xeroxed” or “cloned” at the horizon. One copy would fall through with the infalling observer and the other copy would get scrambled and radiated with the evaporation products. Each observer would see the usual laws of nature respected. The problem is that it potentially violates another law of quantum mechanics. I call it the “Quantum Xerox” principle. It says that quantum information stored in a single system can not be duplicated. To illustrate the principle consider a single spin whose z component can take two values. $\sigma_z = \pm 1$. We call the two states $|+\rangle$ and $|-\rangle$. Suppose we had a quantum xerox machine which could produce a clone of the spin in exactly the original state. Its action is defined by

$$\begin{aligned} |+\rangle &\rightarrow |++\rangle \\ |-\rangle &\rightarrow |--\rangle. \end{aligned} \tag{2.3}$$

But now consider the action of the quantum xerox machine on the superposition of states $|+\rangle + |-\rangle$ which represents a spin oriented along the x axis. According to the principles of quantum mechanics the result must be the superposition $|++\rangle + |--\rangle$. This is because quantum evolution is a linear process. On the other hand the QXM must clone the original state

$$(|+\rangle + |-\rangle) \rightarrow (|+\rangle + |-\rangle) \otimes (|+\rangle + |-\rangle) \tag{2.4}$$

or

$$(|++\rangle + |--\rangle) \rightarrow |++\rangle + |+-\rangle + |-+\rangle + |--\rangle. \tag{2.5}$$

Evidently the quantum Xerox machine is not consistent with the principle of linear quantum evolution.

The Quantum Xerox Principle was probably an implicit part of Stephen’s thinking. I remember that in Aspen sometime around 1992 I gave a seminar explaining why I thought loss of coherence was a big problem and as part of the talk I introduced the *QX* principle. A number of people in the audience didn’t believe it at first but Stephen’s reaction was instantaneous: “So now you agree with me”. I said no, that I didn’t but that I was trying to explain to the high energy people why it was such a serious problem. However, it was also true that I couldn’t see my way out of the paradox.

3 Horizon Complementarity

“When you have eliminated all that is impossible, whatever remains must be the truth, no matter how improbable it is.”

Sherlock Holmes

The principle of Horizon Complementarity I interpret to mean that no observer ever witnesses a violation of the laws of nature. It is obvious that no observer external to the black hole is in danger of seeing the forbidden duplication of infalling information since one copy is behind the horizon. The danger, pointed out by John Preskill [17], is that the observer outside the horizon, call him B , can jump into the horizon having previously collected the relevant information in the Hawking radiation. We must now worry whether the original infalling observer (call her A) can send a signal to B so that B has witnessed information duplication albeit behind the horizon. The answer is no, it is not possible [18]. The reason is interesting and involves a fact first explained by Don Page [19] who realized that to get a single bit of information out of the evaporation products you must wait until about half the entropy of the black hole has been evaporated. This takes a time, in Planck units of order M^3 where M is the black hole mass. It is then easy to see from the black hole metric that if B waits a time of that order of magnitude before jumping behind the horizon, then A must send her signal extremely quickly after passing the Horizon. Otherwise the message will not arrive at B before hitting the singularity. Quantitatively the time that B has after passing the horizon is of order $\Delta t \sim \exp -M^2$ where all things are given in Planck units.

Now in classical physics an arbitrary amount of information can be sent in an arbitrarily small time using an arbitrarily small energy. But in quantum theory if we want to send a single bit we must use at least one quantum. Obviously that quantum must have an energy satisfying $\Delta E \sim \exp M^2$. In other words A had to be carrying an energy vastly larger than the black hole in order that B ever witness the forbidden information cloning. This makes no sense since if A had that much energy it couldn't possibly fit inside the black hole. We see that a consistent use of quantum mechanics simultaneously allows external observations to be consistent with quantum coherence; the infalling observations to be consistent with the equivalence principle; and finally, prevents the observer who chooses to jump in after accumulating some information from ever seeing information duplication.

I can now state the principle of Horizon Complementarity. All it says is that no observer ever sees a violation of the laws of nature. More specifically it says that to an observer

who never crosses the horizon, the horizon behaves like a conventional complex system which can absorb thermalize and re-emit all information that falls on it. No information is ever lost. In essence, the world on the outside of a horizon is a closed system.

On the other hand a freely falling observer encounters nothing out of the ordinary, no large tidal force, high temperature or brick wall at the horizon. The paradox of information being at two places at the same time are apparent and a careful analysis shows that no real contradictions arise.

But there is a weirdness to it. Just to make the point let's imagine a whole galaxy falling into a huge black hole with a Schwarzschild radius equal to a billion light years. From the outside, the galaxy and all its unfortunate inhabitants appear to be heated to Planckian temperature, thermalized and eventually emitted as evaporation products and all this takes place at the horizon! On the other hand, the infalling galactic inhabitants glide through perfectly happily. To them the trauma only happens at the singularity a billion years later. But as in the case of certain life-after-death theories, the folks on the other side can never communicate with us.

It is clear from Horizon Complementarity that a revision is needed in the way we think about information being localized in space-time. In both classical relativity and in quantum field theory the space-time location of an event is invariant, that is, independent of the observer. Nothing in either theory prepares us for the kind of weirdness I described above.

4 The Holographic Principle

The idea that information is in some sense stored at the boundary of a system instead of the bulk was called Dimensional Reduction by 't Hooft and the Holographic Principle by me [4, 5]. The simple argument for the Holographic Principle goes as follows. Imagine a ball of space Γ bounded by a spherical area $A = 4\pi R^2$. Ordinarily we would assume that the entropy in that region can be arbitrarily large. However if we introduce a cutoff in space, say at the Planck scale and assume that within each Planckian volume only a small number of states are possible then it is natural to assume that the maximum entropy is finite and proportional to the volume of the region. However it is easy to see that this can not be so. Imagine an imploding spherical light-like shell with just enough energy so that the boundary of Γ gets turned into a black hole horizon. The final entropy of the system is the Bekenstein-Hawking entropy of the black hole $S_{BH} = A/4G$. By the Second Law the

entropy within Γ could not have been larger than this. But by causality this bound can't depend on the existence of the infalling light like shell. Thus it follows that the largest entropy a region of space can have is given by

$$S_{max} = A/4G. \tag{4.1}$$

A further implication is that the number of degrees of freedom (binary bits) needed to describe a region is also proportional to the area. It is as though the information in the bulk of Γ can be mapped to a set of ‘‘Holographic’’ degrees of freedom on the boundary. The idea that the quantum mechanics of a region can be described in terms of a theory of no more than $A/4G$ degrees of freedom is the content of the Holographic Principle. It is also very weird but it is forced on us by Bekenstein’s observation about black hole entropy [20], the usual interpretation of entropy as counting states and Holmes’ dictum.

5 The Ultraviolet/Infrared Connection

The dominant paradigm of 20th century physics has been that small distance means high energy or more precisely high momentum. Thus to probe increasingly small (and presumably increasingly fundamental) objects, higher and higher energy accelerators have been built. Again this is a lesson learned from conventional quantum field theory. But the more we learn about the combination of gravity and quantum mechanics the clearer it becomes that this trend will eventually reverse itself. Imagine trying to probe distance scales very much smaller than the Planck scale by building the *Gedankatron*, a collider of colossal proportions, which can accelerate electrons to momenta way above the Planck scale. Well, there is no need to build it. We already know what will happen. In a head-on collision a black hole will form and the black hole will have a Schwarzschild radius of order the center of mass energy M in Planck units. The black hole will evaporate with a temperature of order $1/M$. Thus the collision will result in the emission of longer and longer wave length quanta and will probe distance scales of order MG and not $1/M$. This is the simplest example of the UV/IR connection. Similar things happen in perturbative string theory where increasing energy again results in decreasing spatial resolution.

In order to clarify the mechanisms behind the UV/IR connection and show its connection with the Holographic Principle I will use the famous example of the AdS/CFT duality of Maldacena [21]. Let me quickly remind you of this duality. We begin with a stack of N D3-branes in ten dimensional string theory. The branes fill the directions

(1, 2, 3). They are all placed at the origin in the (4, 5, 6, 7, 8, 9) directions. We also define $r^2 = (x^4)^2 + \dots + (x^9)^2$. The geometry of the resulting stack has a horizon at $r = 0$ where the stack is located.

I will not explain in detail the so called decoupling or near-horizon limit. The literature on the subject is enormous. Suffice it to say that the D-brane stack is exactly described in two ways which are dual to one another. The first is by gravity in an AdS space with horizon at $r = 0$. The AdS also has a time-like causal boundary at $r = \infty$. Although the boundary is an infinite proper distance from any finite r , the geometry is such that light takes a finite time to travel to the boundary and back.

The dual ‘‘Holographic’’ description is in terms of a 3+1 dimensional conformal gauge theory whose precise details are unimportant for our purposes. The quantum field theory is usually interpreted as residing on the boundary of the AdS, infinitely far from the horizon. On the other hand the branes that support the open string field quanta are supposedly located at $r = 0$. There seems to be some serious confusion about the location of the branes. Are they at $r = 0$ or $r = \infty$? The answer to this question is closely related to the paradox of where the information in a black hole resides.

In the field theory dual the location of the branes in the $(x^4..x^9)$ space is described by a set of six $N \times N$ matrix valued fields ϕ . The connection between the location of a brane and the corresponding field is

$$\langle r^2 \rangle = \frac{1}{N^2} \langle \text{Tr} \phi^2 \rangle. \quad (5.1)$$

Now we can see the problem in localizing the branes. The field ϕ is a canonically normalized scalar field with mass dimension 1. As such, it has zero point fluctuations which render the value of r^2 divergent. If taken at face value, this would say that the branes are located at the boundary at $r = \infty$. But this is not the most useful way to think about the problem. A better way to think about it is to introduce an ultraviolet cutoff in the gauge theory. Let the regulator frequency be called ν . Then (5.1) is replaced by

$$\langle r^2 \rangle = \frac{1}{N^2} \text{Tr} \langle \phi^2 \rangle = \nu^2. \quad (5.2)$$

Once again we see something weird. The location of the branes is not an invariantly defined quantity. It depends on the frequency resolution that the observer uses. If the branes are observed in a way that averages over high frequency oscillations then their location is near $r = 0$. On the other hand if all the ultraviolet fluctuations are included then the branes are located out near the causal boundary of AdS and the entire theory in the bulk of AdS is described by a holographic description in terms of boundary degrees of

freedom. This interplay between short time cutoffs in the quantum field theory and the location of the brane degrees of freedom is called the UV/IR connection [6].

The UV/IR connection is closely related to the weirdness of Complementarity. Imagine an observer falling toward a black hole horizon while blowing a dog whistle. A dog whistle is just a whistle whose frequency is beyond the range of human hearing. The freely falling observer never hears the whistle. But to someone outside the black hole the frequency is red shifted so that after a while the whistle becomes audible. In the same way, the ultra-high frequency fluctuations of an object are invisible under ordinary circumstances. But as the object approaches the horizon the external observer becomes sensitive to them. In the example I just showed you the observed location and spread of an object depends on the visibility of the high frequency fluctuations. Thus the UV/IR connection provides a mechanism for understanding Horizon Complementarity [22]. Horizon Complementarity the Holographic Principle and the UV/IR connection are different facets of the same weirdness that characterizes the new paradigm.

6 Counting Black Hole Microstates

Ultimately the question of black hole coherence boils down to whether or not the formation and evaporation of a black hole can be described from the outside by an S-matrix. Stephen said no. 't Hooft and I said yes. Although, for years, many string theorists sat on the fence about the issue, they really had no choice. The only mathematically well defined objects in string theory, at least in a flat background, are S-matrix data. This includes the complete list of stable objects in the theory and the transition amplitudes for their scattering. Unstable objects also have meaning as resonances which can be defined as poles of the S-matrix in the complex energy plane. If the laws of quantum mechanics are not violated for a distant observer, black holes are simply such resonances. It sometimes happens that resonances become extremely densely spaced. This occurs in nuclear collisions and also in string theory. In these cases the practical tools are those of statistical mechanics and thermodynamics.

Complementarity implies that the thermodynamics of a black hole should arise from a quantum statistical mechanical origin. Consider the thermodynamic description of a bathtub of hot water. We specify a few macroscopic variables such as the volume, energy temperature etc. Of special significance is the entropy. Entropy, as we know, is a measure of our ignorance of the precise microscopic details of the tub of water. It measures the

logarithm of the number of quantum states consistent with the macroscopic description. The existence of an entropy tells us that there is a hidden set of microscopic degrees of freedom. It doesn't tell us what those degrees of freedom are but it tells us they are there and that they can store detailed information that our thermodynamic description is too coarse grained to see. The principle of Black Hole Complementarity requires the thermodynamics of black holes to originate from the coarse graining of hidden microscopic degrees of freedom and that the entropy is counting the number of microstates of the black hole.

General relativity does not tell us what those micro-states are. But a theory like string theory which is supposed to be a consistent quantum theory of gravity should tell us and it does [7, 8, 9, 10, 11]. Let us begin with a single string in very weakly coupled string theory. If the string is excited to a high state of excitation it typically forms a random walking tangle. It is natural to describe it statistically.

Let L be the total length of string and $T = \frac{1}{\alpha'}$ be the tension. The mass of the string is then

$$m = TL. \tag{6.1}$$

A good model of the string is to think of it as a series of links on a lattice. The link size is the string length $\sqrt{\alpha'}$. Lets suppose that the when we follow the string and it arrives at a lattice site it can continue in n distinct directions. We can then count the number of configurations of the string. The number of links is $L/\sqrt{\alpha'}$ and the number of states is

$$N_{states} = n^{L/\sqrt{\alpha'}}. \tag{6.2}$$

Thus the entropy is

$$S = \frac{L}{\sqrt{\alpha'}} \log n \tag{6.3}$$

or using (6.1) $S = m\sqrt{\alpha'} \log n$. The thing to abstract from this formula is that the entropy is proportional to the mass in string units. The precise coefficient is determined by the details of string theory.

$$S = cm\sqrt{\alpha'}. \tag{6.4}$$

Now imagine turning up the string coupling constant. The large ball of random walking string will begin to experience the effects of gravity. It will shrink and will gain some negative gravitational binding energy. Eventually it will turn into a black hole. This much is obvious. What is less obvious is what happens if we begin with a neutral black hole and turn off the coupling. It is clear that it must evolve into a system of free strings in this

limit but how many such strings will be left at the end of the process? Surprisingly the answer is that the overwhelmingly most likely final state is a single string! The number of configurations of one long string vastly outweighs all other configurations. Thus we can go back and forth from single string states to black holes.

This suggests the following strategy [7]. Start with a black hole of a given mass M . Let us follow it while we adiabatically turn down the string coupling. The entropy will be conserved by such an adiabatic process. At the end we will get a single string of some other mass m . If we can follow the process and determine m we can use (6.4) to estimate the entropy of the original black hole.

This strategy has been used to study a large variety of black holes that occur in string theory [7, 8, 9, 10, 11] including Schwarzschild black holes in all dimensions, and special supersymmetric black holes where the precise coefficients can be obtained. Up to numerical coefficients of order unity, the results always agree with the universal formula

$$S = \frac{area}{4G}. \quad (6.5)$$

By now string theory has given us great confidence in the counting of black hole microstates and that there is no need to look for a new basis for gravitational entropy outside the framework of conventional quantum statistical mechanics.

7 de Sitter Space

Let me turn now to Stephen's favorite subject, cosmology. According to the inflationary theory the universe may have started as a long lived approximation to de Sitter Space with a very small radius of curvature, perhaps a couple of orders of magnitude larger than the Planck radius. Right now observations indicate that it may end as a de Sitter Space of radius $\sim 10^{60}l_p$. Whether or not this will ultimately prove to be the case it clearly behooves us to study and understand the quantum nature of de Sitter Space. Once again, Stephen, together with Gary Gibbons, led the way [23].

de Sitter Space is another example of a space-time that has an event horizon and therefore an entropy and temperature. Accordingly, we should expect paradoxical issues of black hole quantum mechanics to confuse and harass us in this context. The rest of my discussion will concentrate on the quantum mechanics of pure de Sitter Space. However before doing so I will remind you of some facts about Schwarzschild black holes in AdS [24]. You will see why shortly. The Penrose diagram for an AdS-Schwarzschild black hole is a

square bounded on top and bottom by singularities and on the sides by causal boundaries which are at an infinite proper distance. The diagonals of the square are the horizons. Note that the boundary is doubled. As usual, the diagram is divided into 4 regions. Region I is the exterior of the black hole. The Penrose diagram has a boost symmetry which in regions I and III it acts as time translation invariance but with the convention that in III it translates in the negative time-like direction. In regions II and IV it translates in space-like directions.

The AdS black hole has a description in the dual CFT language. It is simply the state of thermal equilibrium for the CFT above the temperature corresponding to the Hawking Page transition [24]. Since the CFT is a conventional quantum system we can be sure that the boundary observers see nothing inconsistent with quantum theory in all possible experiments that they can perform on the black hole. In particular the region I which is in causal contact with the boundary and which is described by the CFT can not experience anything that violates the standard quantum principles. No information can be lost across the horizon. In this respect region I is a closed quantum system.

How does black hole complementarity manifest itself in this system? To see the full implications of exact information conservation in terms of correlation functions is extremely complicated. But Maldacena has given a very useful implication [25]. Consider the correlation function of two local boundary operators at widely separated times t and t' . For example the operators could be the gravitational field evaluated at the boundary and described by the energy-momentum tensor of the QFT. The correlation function can be computed by doing a bulk calculation of the graviton propagator in the AdS-Schwarzschild background. The answer is that it exponentially tends to zero as the time separation grows. From the CFT point of view the exponential decrease has a simple explanation. The thermal environment leads to dissipation which generally causes correlations to disappear exponentially. The coefficient in the exponential is a dissipation coefficient. However, Maldacena argues that there is something wrong with this conclusion. His claim is that the correct answer on general grounds is that the correlation decreases exponentially until it is of order $\exp -S$ where S is the finite black hole entropy. Thereafter it stops decreasing. If Maldacena is correct it means that ordinary QFT in the bulk is missing something important and that something is closely connected to conservation of information outside the black hole. The argument that the correlation has a nonvanishing limit was not spelled out in Maldacena's paper but I will derive it shortly.

Now what does all this have to do with de Sitter Space? To answer this we only

need to draw the Penrose diagram for de Sitter Space. In fact it is identical to the AdS-Schwarzschild case! There are very big geometrical differences but the causal structure is the same. In the de Sitter case the vertical boundaries of the Penrose diagram are not spatial infinity but the “north and south poles” of the spherical spatial sections. One of these poles we can identify as “the observer”. The horizontal boundaries are not high-curvature singularities but are instead the infinitely inflated past I^- and the infinitely inflated future I^+ .

The metric for de Sitter Space is

$$ds^2 = R^2 \left(dt^2 - \cosh^2(t) d\Omega_d^2 \right) \quad (7.1)$$

where R is the radius of the de Sitter Space and d is the dimension of space.

If de Sitter Space exists in a quantum theory of gravity, perhaps the most urgent question is what are the mathematical objects that the theory defines. This is particularly important for string theory. In flat space the answer is S-matrix data. In AdS it is boundary correlators of the gravitational and other bulk fields. In both cases we go to the boundary of the world and define vertex operators whose correlators are the “definables”. The obvious suggestion for de Sitter Space is that we do the same thing. The boundary of de Sitter Space is the union of I^- and I^+ . This suggests that the definables consist of S-matrix-like elements which relate initial states on I^- to final states on I^+ [26, 27, 28]. I have used the term definables ¹ and not observables for the reason that no one can ever observe them. The argument is closely connected to that of Section 3 about why information duplication can’t be detected behind a black hole horizon. In that case there was just not enough time for any observer to collect information from the observers A and B before reaching the singularity. The similarity of I^+ to the singularity of a black hole means that two observers at fixed spatial coordinates can not transmit the results of measurements to one another once global time gets too late. Accordingly correlators of fields at the future boundary are unobservable.

In the black hole case it is not just that quantum correlators on the singularity are un-measurable. Just their mere *mathematical* existence would require us to trace over them in defining final states. It is of obvious importance to know if the same is true in de Sitter Space. In other words can similar arguments show that the existence of de Sitter Space boundary correlators will lead some observer to see a violation of the laws of

¹Witten has used the terms computables and meta-observable for the same objects.

nature? I will argue that the answer is yes. But first we need to formulate an appropriate Complementarity principle.

In the black hole case, a convenient starting point for discussing observations outside the hole is the presentation of the geometry in Schwarzschild coordinates. The distinguishing features of these coordinates are that they are static and that they only cover the region on the observer's side of the horizon. In de Sitter Space an observer means a time-like trajectory that begins on I^- and ends on I^+ . All such pairs of points are related by symmetry and it is always possible to choose them at the same spatial location. Thus we can always choose the observer to be the $r = 0$ edge of the Penrose diagram. By analogy with the AdS/Schwarzschild case the diagonals of the square Penrose diagram are horizons and the region on the observers side of the horizon is a triangle. On this region it is possible to choose static coordinates so that the metric takes the form

$$ds^2 = R^2 \left((1 - r^2) dt^2 - (1 - r^2)^{-1} dr^2 - r^2 d\Omega_{d-1}^2 \right). \quad (7.2)$$

This geometry has a horizon at $r = 1$ which now surrounds the observer. As usual, the horizon has an entropy

$$S = \text{area}/4G = \pi R^2/G \quad (7.3)$$

and a temperature. The proper temperature at a point r is given by

$$T(r) = \frac{1}{2\pi R \sqrt{1 - r^2}} \quad (7.4)$$

As for Rindler or Schwarzschild space, the temperature diverges near the horizon. The observer at $r = 0$ experiences a more modest temperature.

The statement of Horizon Complementarity is fairly obvious. An observer at $r = 0$ sees the world as a closed finite system at a non-zero temperature. Once again, closed means that no information is lost and the system obeys the rules of quantum mechanics for such closed systems [29]. What are the implications of the complementarity principle and especially the conservation of information? As in the similar AdS/Schwarzschild case there are implications for the long time behavior of correlators. However before discussing them I want to consider the behavior of field correlators in ordinary quantum field theory in the fixed de Sitter Space background. For this purpose I return to global coordinates.

For simplicity I will use the example of a massive scalar field Φ with mass μ/R . The field equation is easily worked out and the asymptotic behavior of Φ is seen to be [26]

$$\Phi \rightarrow \exp -\gamma|t| \quad (7.5)$$

as $t \rightarrow \pm\infty$. The “diffusion constant” γ is given by

$$\gamma = d \pm i\sqrt{\mu^2 - \frac{d^2}{4}}. \quad (7.6)$$

The complex value of γ has a simple meaning in terms of damped oscillations. Depending of the value of μ the oscillations of the field are either under-damped or over-damped. The real part of γ is always positive.

Now let us return to static coordinates and consider the correlator at $r = 0$ and large time separation. Time translation invariance in the static patch together with (7.5) requires the correlator to behave like

$$F(t) = \langle \phi(t)\phi(t') \rangle \sim \exp -\gamma|t - t'|. \quad (7.7)$$

Thus, as in the AdS/Schwarzschild black hole the correlator evaluated by naive quantum field theory tends to zero exponentially with time. As we will see, this is inconsistent with the finite entropy of the static patch.

8 Correlations in Finite Entropy Systems

Let us consider an arbitrary system described by Hamiltonian H at temperature $1/\beta$. I will not assume anything about the total number of states of the system but only that the thermal entropy is finite, $S = \text{finite}$. An immediate implication is that the spectrum is discrete and that the level spacing is of order

$$\Delta E \sim \exp -S. \quad (8.1)$$

Let's label the energy levels E_m .

Now consider a correlator of the form [30]

$$\langle \Phi(0)\Phi(t) \rangle \equiv F(t). \quad (8.2)$$

It is defined by

$$\begin{aligned} F(t) &= \frac{1}{Z} \text{Tr} e^{-\beta H} \Phi(0)\Phi(t) \\ &= \sum_{mn} e^{-\beta E_n} |\Phi_{nm}|^2 e^{it(E_m - E_n)}. \end{aligned} \quad (8.3)$$

For simplicity I will assume that the diagonal matrix elements of Φ vanish. This implies that the time average of Φ also vanishes.

I want to determine if F tends to zero as $t \rightarrow \infty$. A simple way to do this is to compute the long-time average of F^*F .

$$L = \lim_{T \rightarrow \infty} \frac{1}{2T} \int_{-T}^T F^*(t) F(t). \quad (8.4)$$

If $F \rightarrow 0$ then $L = 0$.

Using (8.3) it is straightforward to compute L ;

$$L = \frac{1}{Z^2} \sum_{mnr s} e^{-\beta(E_m + E_r)} |\Phi_{mn}|^2 |\Phi_{rs}|^2. \quad (8.5)$$

This is a positive definite quantity. We have therefore proved that the correlation function does not go to zero.

It is not hard to estimate L [31]. We use the fact that the level spacing is of order $\exp -S$ which implies that the matrix elements of Φ are of the same order of magnitude. The result is

$$L \sim \exp -2S \quad (8.6)$$

indicating that at asymptotic times the average magnitude of the correlator is of order

$$|F| \sim \exp -S \quad (8.7)$$

which agrees with Maldacena's guess.

However, this does not mean that the correlator tends smoothly to an exponentially small constant. In fact it does something quite different. It becomes “noisy”. Typically the fluctuations are small, of order $\exp -S$. But if you wait long enough large fluctuations occur. On sufficiently long time scales the correlation function return to close to its original value. In [30] the reader can find plots of the results of some numerical studies done by Lisa Dyson and James Lindesay [30] which show the typical behavior. The large scale fluctuations are the quantum version of Poincare recurrences in classical mechanics. They are just an indication that in a closed finite system, if you wait long enough everything will eventually happen and not just once but in eternal repetition.

What is the significance of this fact and why is it missed by quantum field theory in the de Sitter Space background? First of all it means that any formulation of quantum gravity in de Sitter Space that relies on the existence of well defined asymptotic fields must fail. Among these I include the possibility that string theory might provide a set of scattering amplitudes relating asymptotic states on $I^\pm$. Another version of de Sitter Space quantum

gravity is the so called dS/CFT correspondence inspired by the AdS/CFT duality. This theory attempts to organize boundary correlators into an Euclidean quantum field theory.

A question which often comes up in any discussion of de Sitter Space is how the static patch observer should think about objects (conventionally called elephants) which enter and leave the static patch through the diagonal past and future horizons. Such events are described by very rare fluctuations which materialize among the large number of low frequency degrees of freedom very near the horizon. In other words they are the very intermittent large fluctuations that can be seen in the numerical simulations of [30]. Such elephants can not be tossed in from I^- . To understand why, let's think about the elephant in its own frame. Because of the thermal fluctuations in de Sitter Space an elephant will eventually be thermalized and turned into black body radiation. Since I^- is an infinite proper time in the past any elephants that were present in the initial conditions have long been thermalized by any finite time. Elephants in de Sitter Space are thermal fluctuations which form by chance and then evaporate.

Finally, why does quantum field theory in the de Sitter Space background miss these fluctuations? The answer is simply that in quantum field theory the entropy of a horizon has an ultraviolet divergence due to modes arbitrarily close to the horizon. These modes provide an infinite number of low frequency oscillators which effectively make the system open. But we know that most of these modes must be eliminated in a proper theory of quantum gravity since the entropy of horizons is finite.

If boundary correlators and S-matrix elements can not exist in de Sitter Space what then are we to base a quantum theory on? Here things are much more obscure than in AdS. A possible answer is to define a theory in the closed static patch. But, as emphasized by Bohr, the clearest use of quantum mechanics is when we can separate the world into a system and a large or classical apparatus (observer). That is just what we can not do in a closed finite thermal system like de Sitter Space. In fact the whole discussion of static patches being defined by infinitely long lived observers in de Sitter Space is internally inconsistent. Like elephants, any real observer is subject to bombardment by the thermal radiation and will become degraded. Exactly how to think about quantum mechanics is far from clear. Perhaps eternal de Sitter Space is just a bad idea and will go away. Perhaps not.

So you can see that Stephen and I have had much to disagree about. These disagreements have been incredibly exciting and certainly provided the high point of my own career. There is no doubt that Stephen's remarkable question about black holes has left

an indelible mark on me but more importantly, on the future of physics.

Happy birthday, Stephen and best wishes for many more.

References

- [1] Stephen Hawking, Breakdown Of Predictability In Gravitational Collapse, Phys.Rev. D14 (1976) 2460 Hawking
- [2] L. Susskind, L. Thorlacius, J. Uglum, The Stretched Horizon and Black Hole Complementarity, hep-th/9306069, Phys. Rev. D48 (1993) 3743-3761
- [3] C.R. Stephens, G. 't Hooft, B.F. Whiting, Black Hole Evaporation without Information Loss, gr-qc/9310006, Class.Quant.Grav. 11 (1994) 621-648
- [4] G. 't Hooft, Dimensional Reduction in Quantum Gravity, gr-qc/9310026
- [5] L. Susskind, The World as a Hologram, J.Math.Phys. 36 (1995) 6377-6396 hep-th/9409089
- [6] L. Susskind, Edward Witten, The Holographic Bound in Anti-de Sitter Space, hep-th/9805114
- [7] Leonard Susskind, Some Speculations about Black Hole Entropy in String Theory, hep-th/9309145
- [8] Ashoke Sen, Black Hole Solutions in Heterotic String Theory on a Torus, Nucl.Phys. B440 (1995) 421-440, hep-th/9411187
- [9] A. Strominger, C. Vafa, Microscopic Origin of the Bekenstein-Hawking Entropy, Phys.Lett. B379 (1996) 99-104, hep-th/9601029
- [10] Curtis G. Callan, Juan M. Maldacena, D-brane Approach to Black Hole Quantum Mechanics, Nucl.Phys. B472 (1996) 591-610, hep-th/9602043
- [11] Gary T. Horowitz, Joseph Polchinski, A CORRESPONDENCE PRINCIPLE FOR BLACK HOLES AND STRINGS. Phys.Rev.D55:6189-6197,1997, hep-th/9612146
- [12] G. 't Hooft, On The Quantum Structure Of A Black Hole, Nucl.Phys. B256 (1985) 727

- [13] Mark Srednicki, Entropy and Area, Phys.Rev.Lett. 71 (1993) 666, hep-th/9303048
- [14] For an early critique of Hawking’s proposal, see D. N. Page, Phys. Rev. Lett. 44 (80) 301.
- [15] T. Banks, M. Peskin, L. Susskind, Difficulties For The Evolution Of Pure States Into Mixed States, Nucl.Phys. B244 (1984) 125
- [16] G. ’t Hooft, The Black Hole Interpretation of String Theory, Nucl. Phys. B335 (90) 138, and references therein.
- [17] John Preskill, Unpublished, 1993
- [18] L. Susskind, L. Thorlacius, Gedanken Experiments involving Black Holes, Phys.Rev. D49 (1994) 966-974, hep-th/9308100
- [19] Don N. Page, Average Entropy of a Subsystem, Phys.Rev.Lett. 71 (1993) 1291-1294, gr-qc/9305007
- [20] J.D. Bekenstein, “Black Holes And Entropy” Phys. Rev. D7, 2333 (1973)
- [21] J. Maldacena, “The Large N Limit of Superconformal Field Theories and Supergravity,” Adv. Theor. Math. Phys. **2**, 231 (1998) [hep-th/9711200].
- [22] L. Susskind, String Theory and the Principle of Black Hole Complementarity, Phys.Rev.Lett. 71 (1993) 2367-2368, hep-th/9307168
- [23] G.W. Gibbons, S.W. Hawking, COSMOLOGICAL EVENT HORIZONS, THERMODYNAMICS, AND PARTICLE CREATION, Phys.Rev.D15:2738-2751,1977
- [24] S.W. Hawking, Don N. Page, THERMODYNAMICS OF BLACK HOLES IN ANTI-DE SITTER SPACE Commun.Math.Phys.87:577,1983
- [25] Juan M. Maldacena, Eternal Black Holes in AdS, hep-th/0106112
- [26] Raphael Bousso, Alexander Maloney, Andrew Strominger, Conformal Vacua and Entropy in de Sitter Space, hep-th/0112218 and references therein.
- [27] E. Witten, Quantum Gravity In De Sitter Space, hep-th/0106109

- [28] Marcus Spradlin, Anastasia Volovich, Vacuum States and the S-Matrix in dS/CFT, hep-th/0112223
- [29] Complementarity in the context of deSitter Space is also discussed in T.Banks, W.Fischler, M-theory observables for cosmological space-times, hep-th/0102077
- [30] Lisa Dyson, James Lindesay, Leonard Susskind, Is There Really a de Sitter/CFT Duality, hep-th/0202163
- [31] Mark Srednicki, The approach to thermal equilibrium in quantized chaotic systems, J. Phys. A 32 (1999) 1163, cond-mat/9809360